Connecticut Maternal Mortality Review Program

Screening Tool to Identify Social Determinants of Health and Discrimination Contributing to Maternal Death

Check	factors that have been identified through record review:		From the record discrimination to Discrimination retreatment of differences on the differences on the differences of the differ	pased on each of fers to unjust or grent categories of	f these factors? prejudicial of people,
Detient	Socio-Demographic Factors		YES	NO	UNK
Patient	race or ethnicity		15	NO 🗆	UNK
	age (< 20 y)			-H	
H	income (Medicaid as a proxy)		H	-H $-$	H
	immigration or citizenship status		H	H	H
H	education (high school degree or less)		H	H	H
	sexual orientation or gender identity				
	marital status (never married)		Ħ	H	H
	weight (obesity)		Ħ	Ħ	Ħ
H	language		Ħ	Ħ	Ħ
	substance abuse				
	other (please specify)		Ī	\Box	Π
	1 77		_	_	_
Check	factors that have been identified through record review:				
Patient	Health Knowledge, Beliefs, Behaviors	Relatio	nal & Psychosocial Fa	ctors	
	functional or health literacy		childhood abuse or tra		
	substance use during pregnancy		mental health condition	ns	
	late or no prenatal care		intimate partner violer	ice	
	lack of compliance with medical treatment		current, past, or pend	ing DCF involver	ment
	leaving against medical advice		lack of social support	or social isolation	n
	repeated ED visits for urgent care		food insecurity		
	refusal of social service referrals		housing insecurity or	nstability	
	other (please specify)		other (please specify)		
Comm	unity & Neighborhood Factors	Structu	ıral Factors		
	lack of community supports (home visiting, case mgt)		lack of access to heal		
	crime	Щ.	lack of adequate heal	th insurance cov	erage
	poverty		other (please specify)		
Щ	environmental exposure				
\sqcup	other (please specify)				
1					

Rev. 07/30/2020

Connecticut Maternal Mortality Review Program

Screening Tool to Identify Social Determinants of Health and Discrimination Contributing to Maternal Death

Check	factor	rs that have been identified through record review:				
Provid	der & I	Facility Factors	Check tl	ne level at which	the factor	was observed:
	cult	ural incompetence (lack of translator, patient education				
		erials, diversity in signage, magazines, posters)		provider		facility policies / practices
		of ongoing training on implicit & explicit bias, structural &		nun ida u		facility maliains / munchines
Ш		rpersonal racism, shared decision-making ative patient-provider/facility interactions (stigmatizing		provider		facility policies / practices
		guage, dismissing concerns, lack of respect, conflict)		provider		facility policies / practices
		essive gatekeeping (inability to reach provider, leaving		providor		radiity poiloide / pradado
		ssages, etc.)		provider		facility policies / practices
	indi	cated labs not ordered or delay in lab work ordered				facility policies / practices
	11			provider		for the contract of the contra
Ш	lack	of follow-up with patient	Ш	provider	Ш	facility policies / practices
	othe	er (please specify)				
		Decisions & Recommendations	Check tl	ne level at which	the factor	was observed:
	sisten	t with Best Practices	Check tl		the factor	
	sisten a)	t with Best Practices over-treatment	Check tl	provider	the factor	facility policies / practices
	sisten a) b)	t with Best Practices over-treatment under-treatment	Check tl	provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c)	t with Best Practices over-treatment under-treatment delay in treatment	Check th	provider	the factor	facility policies / practices facility policies / practices facility policies / practices
	sisten a) b)	t with Best Practices over-treatment under-treatment	Check th	provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists	Check th	provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning	Check th	provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum	Check th	provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals	Check th	provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h) i)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals lack of screening for depression and intimate partner violence	Check th	provider provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals	Check th	provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h) i)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals lack of screening for depression and intimate partner violence	Check th	provider provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h) i)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals lack of screening for depression and intimate partner violence	Check th	provider provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h) i)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals lack of screening for depression and intimate partner violence	Check th	provider provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h) i)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals lack of screening for depression and intimate partner violence	Check th	provider provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h) i)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals lack of screening for depression and intimate partner violence	Check th	provider provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h) i)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals lack of screening for depression and intimate partner violence	Check th	provider provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h) i)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals lack of screening for depression and intimate partner violence	Check th	provider provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h) i)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals lack of screening for depression and intimate partner violence	Check th	provider provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h) i)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals lack of screening for depression and intimate partner violence	Check th	provider provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h) i)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals lack of screening for depression and intimate partner violence	Check th	provider provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices
	a) b) c) d) e) f) g) h) i)	t with Best Practices over-treatment under-treatment delay in treatment inadequate pain management assumptions about patient adherence to treatment lack of, or delay in, referrals to specialists inadequate discharge planning inadequate care-coordination prenatally and postpartum lack of, or delay in, social service referrals lack of screening for depression and intimate partner violence	Check th	provider provider provider provider provider provider provider provider provider provider	the factor	facility policies / practices facility policies / practices

Rev. 07/30/2020 2